

Aryabhatta Knowledge University (AKU)

Information Technology (IT)

Mathematics-III

Solved Exam Paper 2019

**Q 1 a) If  $y=A \cos(\log x) + B \sin(\log x)$ , Show that  $x^2y_{n+2} + (2n+1) * y_{n+1} + (n^2 + 1) y_n = 0$  where  $y_n = \frac{d^n y}{dx^n}$**

If  $y = A\cos(\log x) + B\sin(\log x)$  ----- (1)

Differentiating (1) w.r.t x, we get

$$y_1 = -A\sin(\log x)/x + B\cos(\log x)/x$$

$$xy_1 = -A\sin(\log x) + B\cos(\log x) \text{ ----- (2)}$$

Diff 2 again w.r.t x, then we get

$$xy_2 + y_1 = -A\cos(\log x)/x - B\sin(\log x)/x$$

$$x^2y_2 + xy_1 = -[\cos(\log x) + \sin(\log x)]$$

$$x^2y_2 + xy_1 = -y$$

$$y_2x^2 + y_1x + y = 0 \text{ ----- (3)}$$

Diff 3 by Leibnitz theorem n times, we get

$$[y_{n+2}x^2 + nc1 y_{n+1}2x + nc2 y_n \cdot 2] + [y_{n+1}x + nc1 y_{n-1}] + y_n = 0$$

$$x^2 y_{n+2} + 2nxy_{n+1} + xy_{n+1} + n(n-1)y_n + ny_n + y_n = 0$$

$$x^2 y_{n+2} + (2n+1)xy_{n+1} + (n^2+1)y_n = 0$$

**Q2 a) Show that the following function is continuous at the point (0,0):**

$$f(x,y) = \begin{cases} \frac{2x^3 + 3y^3}{x^2 + y^2}, & (x,y) \neq (0,0) \\ 0, & (x,y) = (0,0) \end{cases}$$

Answer:

$$f(x,y) = \begin{cases} \frac{2x^3 + 3y^3}{x^2 + y^2}, & (x,y) \neq (0,0) \end{cases}$$

$$0, \quad (x,y) = (0,0)$$

$$0 \leq \left| \frac{2x^3 + 3y^3}{x^2 + y^2} \right| \leq \frac{2|x^3|}{x^2 + y^2} + \frac{3|y^3|}{x^2 + y^2} = 0$$

$$0 \leq \frac{2|x^3|}{x^2} + \frac{3|y^3|}{y^2} = 2|x| + 3|y|$$

$$\lim_{(x,y) \rightarrow (0,0)} 2|x| + 3|y| = 0$$

∴ by the squeeze theorem, we conclude that

$$\lim_{(x,y) \rightarrow (0,0)} f(x,y) = (0,0) \text{ that is } \Rightarrow \lim_{(x,y) \rightarrow (0,0)} \frac{2x^3 + 3y^3}{x^2 + y^2} = 0$$

∴  $f$  is continuous at  $(0,0)$

**Q2 b) If  $z(x+y) = x^2 + y^2$  show that**

$$\left(\frac{\partial z}{\partial x} - \frac{\partial z}{\partial y}\right)^2 = 4\left(1 - \frac{\partial z}{\partial x} - \frac{\partial z}{\partial y}\right)$$

solution:

$$z(x+y) = x^2 + y^2$$

$$z = \frac{x^2 + y^2}{x+y}$$

$$\frac{\partial z}{\partial x} = \frac{(x+y)2x - (x^2 + y^2).1}{(x+y)^2} = \frac{x^2 + 2xy - y^2}{(x+y)^2}$$

$$\frac{\partial z}{\partial y} = \frac{(x+y)2y - (x^2 + y^2).1}{(x+y)^2} = \frac{-x^2 + 2xy + y^2}{(x+y)^2}$$

$$(\frac{\partial z}{\partial x} - \frac{\partial z}{\partial y}) = \frac{x^2 + 2xy - y^2}{(x+y)^2} = \frac{-x^2 + 2xy + y^2}{(x+y)^2} = \frac{2(x-y)}{(x+y)}$$

$$(\frac{\partial z}{\partial x} - \frac{\partial z}{\partial y})^2 = \frac{4(x-y)^2}{(x+y)^2}$$

-----1

$$4(1 - \frac{\partial z}{\partial x} - \frac{\partial z}{\partial y}) = 4 [ 1 - \frac{x^2 + 2xy - y^2}{(x+y)^2} - \frac{-x^2 + 2xy + y^2}{(x+y)^2} ]$$

$$= \frac{4(x-y)^2}{(x+y)^2}$$

-----2

From 1 & 2

$$(\frac{\partial z}{\partial x} - \frac{\partial z}{\partial y})^2 = 4(1 - \frac{\partial z}{\partial x} - \frac{\partial z}{\partial y})$$

### Q3 a) Transform the equation

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0 \text{ into polar coordinates.}$$

Solution:

we have  $x = r\cos\theta$  ,  $y = r\sin\theta$

$$r^2 = x^2 + y^2 , \theta = \tan^{-1}y/x$$

$$\frac{\partial u}{\partial x} = \frac{\partial u}{\partial r} \cdot \frac{\partial r}{\partial x} + \frac{\partial u}{\partial \theta} \cdot \frac{\partial \theta}{\partial x} \Rightarrow \frac{\partial u}{\partial x} (\cos \theta) - \frac{\partial u \sin \theta}{\partial \theta r}$$

$$\frac{\partial^2 u}{\partial x^2} = \frac{\partial}{\partial x} \left( \frac{\partial u}{\partial x} \right)$$

$$\begin{aligned}
&= \left( \cos\theta \frac{\partial}{\partial r} - \frac{\sin\theta}{r} \frac{\partial}{\partial \theta} \right) \left( \cos\theta \frac{\partial u}{\partial r} - \frac{\sin\theta \partial u}{r \partial \theta} \right) \\
&= \cos\theta \frac{\partial}{\partial x} \left( \cos\theta \frac{\partial u}{\partial r} - \frac{\sin\theta \partial u}{r \partial \theta} \right) - \frac{\sin\theta}{r} \frac{\partial}{\partial \theta} \left( \cos\theta \frac{\partial u}{\partial r} - \frac{\sin\theta \partial u}{r \partial \theta} \right) \\
&= \cos\theta \left( \cos\theta \frac{\partial^2 u}{\partial r^2} + \frac{\sin\theta \partial u}{r^2 \partial \theta} - \frac{\sin\theta \cdot \frac{\partial^2 u}{\partial r \partial \theta}}{r} \right) - \frac{\sin\theta}{r} \left( -\sin\theta \frac{\partial u}{\partial r} + \right. \\
&\quad \left. \cos\theta \frac{\partial^2 u}{\partial r \partial \theta} - \frac{\cos\theta \partial u}{r \partial \theta} - \frac{\sin\theta \partial^2 u}{r \partial \theta^2} \right) \\
&= \cos^2 \theta \frac{\partial^2 u}{\partial r^2} + \frac{\sin\theta \cos\theta \partial u}{r^2 \partial \theta} - \frac{\sin\theta \cos\theta \frac{\partial^2 u}{\partial r \partial \theta}}{r} + \frac{\sin^2 \theta \partial u}{r \partial r} -
\end{aligned}$$

$$\begin{aligned}
&\frac{\sin\theta \cos\theta}{r} \frac{\partial^2 u}{\partial \theta \partial u} + \frac{\sin\theta \cos\theta \partial u}{r^2 \partial \theta} + \frac{\sin^2 \theta \partial^2 u}{r^2 \partial \theta^2} \\
&= \cos^2 \theta \frac{\partial^2 u}{\partial r^2} + 2 \frac{\sin\theta \cos\theta \partial u}{r^2 \partial \theta} - 2 \frac{\sin\theta \cos\theta \frac{\partial^2 u}{\partial r \partial \theta}}{r} + \frac{\sin^2 \theta \partial u}{r \partial r} +
\end{aligned}$$

$$\frac{\sin^2 \theta \partial^2 u}{r^2 \partial \theta^2}$$

(1)

$$\begin{aligned}
&= \frac{\partial u}{\partial y} = \frac{\partial u}{\partial r} \frac{\partial r}{\partial y} + \frac{\partial u}{\partial \theta} \frac{\partial \theta}{\partial y} \\
&= \frac{\partial u}{\partial r} y/r + \frac{\partial u}{\partial \theta} \frac{x}{(x^2 + y^2)} \\
&= \frac{\partial u}{\partial r} \sin\theta + \frac{\partial u}{\partial \theta} \frac{\cos\theta}{r}
\end{aligned}$$

$$= \frac{\partial^2 u}{\partial y^2} = \frac{\partial}{\partial y} \left( \frac{\partial u}{\partial y} \right)$$

$$= (\sin\theta \frac{\partial}{\partial u} + \frac{\cos\theta \partial}{r \partial \theta}) (\sin\theta \frac{\partial u}{\partial r} + \frac{\cos\theta \partial u}{r \partial \theta})$$

$$= \sin\theta \frac{\partial}{\partial r} (\sin\theta \frac{\partial u}{\partial r} + \frac{\cos\theta \partial u}{r \partial \theta}) + \frac{\cos\theta \partial}{r \partial \theta} (\sin\theta \frac{\partial u}{\partial r} + \frac{\cos\theta \partial u}{r \partial \theta})$$

$$= \sin\theta \left[ \sin\theta \frac{\partial^2 u}{\partial r^2} - \frac{\cos\theta \partial u}{r^2 \partial \theta} + \frac{\cos\theta}{r} + \frac{\partial^2 u}{\partial r \partial \theta} \right] + \frac{\cos\theta}{r} [\cos\theta \left[ \frac{\partial u}{\partial r} + \right.$$

$$\left. \sin\theta \frac{\partial^2 u}{\partial r \partial \theta} - \frac{\sin\theta \partial u}{r \partial \theta} + \frac{\cos\theta \partial^2 u}{r \partial \theta^2} \right]$$

$$\begin{aligned}
 &= \sin^2\theta \frac{\partial^2 u}{\partial r^2} - \frac{\sin\theta \cos\theta \partial u}{r^2} + \frac{\sin\theta \cos\theta}{r} \frac{\partial^2 u}{\partial r \partial \theta} + \frac{\cos^2\theta}{r} \left( \frac{\partial u}{\partial r} \right) + \frac{\sin\theta \cos\theta}{r} \\
 &= \sin^2\theta \frac{\partial^2 u}{\partial r^2} - 2 \frac{\sin\theta \cos\theta \partial u}{r^2} + 2 \frac{\sin\theta \cos\theta}{r} \frac{\partial^2 u}{\partial r \partial \theta} + \frac{\cos^2\theta}{r} \left( \frac{\partial u}{\partial r} \right) + \\
 &\quad \frac{\frac{\partial^2 u}{\partial \theta^2}}{\dots} \\
 (2)
 \end{aligned}$$

By adding 1 & 2

$$\begin{aligned}
 \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} &= (\sin^2\theta + \cos^2\theta) \frac{\partial^2 u}{\partial r^2} + (\sin^2\theta + \cos^2\theta) 1/r \frac{\partial u}{\partial r} + (\sin^2\theta + \cos^2\theta) 1/r^2 \frac{\partial^2 u}{\partial \theta^2} \\
 &= \left( \frac{\partial^2 u}{\partial r^2} + 1/r \frac{\partial u}{\partial r} + 1/r^2 \frac{\partial^2 u}{\partial \theta^2} \right) \text{ ans}
 \end{aligned}$$

**Q4 Find the extreme values of  $f(x,y,z) = 2x+3y+z$ , such that  $x^2+y^2=5$  and  $x+z=1$**

$$f(x,y,z) = 2x + 3y + z \quad \dots \quad (1)$$

$$f(x,y) = (x^2 + y^2) - 5 \quad \dots \quad (2)$$

$$y(x,z) = x+z-1 \quad \dots \quad (3)$$

Lagranges Multipliers Equations are

$$\frac{\partial f}{\partial x} + \lambda \frac{\partial \phi}{\partial x} + m \frac{\partial \psi}{\partial x} = 0$$

$$\frac{\partial f}{\partial y} + \lambda \frac{\partial \phi}{\partial y} + m \frac{\partial \psi}{\partial y} = 0$$

$$\frac{\partial f}{\partial z} + \lambda \frac{\partial \phi}{\partial z} + m \frac{\partial \psi}{\partial z} = 0$$

$$2 + \lambda(2x) + m(1) = 0 \quad \dots \dots \dots (4)$$

$$3 + \lambda(2y) + m(0) = 0 \quad \dots \dots \dots (5)$$

$$1 + \lambda(0) + m(1) = 0 \quad \dots \dots \dots (6) \Rightarrow m = -1$$

putting the values of m in (4) and (5), we get

$$2 + 2\lambda x - 1 = 0 \Rightarrow 2\lambda x = -1, \quad x = -\frac{1}{2\lambda}$$

$$3 + 2\lambda y = 0 \Rightarrow 2\lambda y = -3, \quad y = -\frac{3}{2\lambda}$$

putting the values of x,y in  $x^2 + y^2 = 5$ , we get

$$\frac{1}{4\lambda^2} + \frac{9}{4\lambda^2} = 5 \Rightarrow \frac{10}{4\lambda^2} = 5$$

$$2\lambda^2 = 1 \Rightarrow \lambda = \pm \frac{1}{\sqrt{2}}$$

$$\text{we know that, } x = -\frac{1}{2\lambda} = \pm \frac{1}{\sqrt{2}} = \pm \frac{1}{\sqrt{2}}$$

$$y = -\frac{3}{2\lambda} = \pm \frac{3\sqrt{2}}{2} = \pm \frac{3}{\sqrt{2}}$$

From (3),  $x+z=1$  or  $z=1-x$

$$z = 1 \pm \frac{1}{\sqrt{2}}$$

putting  $x = \frac{1}{\sqrt{2}}$ ,  $y = \frac{3}{\sqrt{2}}$  and  $z = 1 - \frac{1}{\sqrt{2}}$  in equation (1), we get

$$f = \frac{2}{\sqrt{2}} + \frac{9}{\sqrt{2}} + 1 - \frac{1}{\sqrt{2}} = \frac{10}{\sqrt{2}} + 1 = 5\sqrt{2} + 1$$

putting  $x = -\frac{1}{\sqrt{2}}$ ,  $y = -\frac{3}{\sqrt{2}}$  and  $z = 1 + \frac{1}{\sqrt{2}}$  in equation (1), we get

$$f = 2\frac{1}{\sqrt{2}} + 3\left(\frac{-3}{\sqrt{2}}\right) + \left(1 + \frac{1}{\sqrt{2}}\right) = -\frac{2}{\sqrt{2}} - \frac{9}{\sqrt{2}} + 1 + \frac{1}{\sqrt{2}}$$

**Q5 Evaluate  $\oint\limits_S \mathbf{F} \cdot d\mathbf{s}$  where  $\mathbf{F} = 4xz \mathbf{i} + y^2 \mathbf{j} + yz \mathbf{k}$  a surface of the cube bounded by  $x=0, y=1, z=0$  , , using Gauss divergence theorem**

s.no

surface

1

OABC

 $-\mathbf{k}$  $dxdy$ 

z

2

DEFG

 $\mathbf{k}$  $dxdy$ 

z

3

OAFG

 $-\mathbf{j}$  $dxdz$ 

y

4

BCDE

 $\mathbf{j}$  $dxdz$ 

y

5

ABEF

 $\mathbf{i}$  $dydz$ 

x

6

OCDG

 $-\mathbf{i}$  $dydz$ 

x

$$\oint\limits_S \mathbf{F} \cdot \mathbf{n} ds = \oint\limits_{OABC} \mathbf{F} \cdot \mathbf{n} ds + \oint\limits_{DEFG} \mathbf{F} \cdot \mathbf{n} ds + \oint\limits_{OAFG} \mathbf{F} \cdot \mathbf{n} ds$$

$$\oint_{BCDE} F^- \cdot n^+ ds + \oint_{ABEF} F^- \cdot n^+ ds +$$

$$\oint_{OCDG} F^- \cdot n^+ ds$$

----- (

$$\oint_{OABC} F^- \cdot n^+ ds = \oint_{OABC} (4xz i^+ + y^2 j^+ + yzk^+) (-k) dx dy$$

$$= \int_0^1 \int_0^1 -yz dx dy = 0 \text{ (as } z = 0)$$

$$\oint_{DEFG} (4xz i^+ + y^2 j^+ + yzk^+) (k) dx dy = \oint_{DEFG} yz dx dy$$

$$\int_0^1 \int_0^1 y(1) dx dy = \int_0^1 dx \left[ \frac{y^2}{2} \right]_{0-1} = [x]_{0-1} = 1/2$$

$$\oint_{OAFG} (4xz i^+ + y^2 j^+ + yzk^+) (-j^+) dx dz = \oint_{OAFG} y^2 dx dz = 6 \text{ (as } y = 0)$$

$$\oint_{BCDE} (4xz i^+ + y^2 j^+ + yzk^+) (j^+) dx dz = \oint_{BCDE} (-y^2) dx dz$$

$$-\int_0^1 dx \int_0^1 dz = (x)_{0-1} (z)_{0-1} = -1 \quad (\text{as } y = 1)$$

$$\oint_{ABEF} (4xz i^+ + y^2 j^+ + yzk^+) \cdot i^+ dy dz = \oint 4xz dy dz$$

$$= \int_0^1 \int_0^1 4(1) z dy dz \Rightarrow 4(y)_{0-1} \left( \frac{z^2}{2} \right)_{0-1} = 4(1) \left( \frac{1}{2} \right) = 2$$

$$\oint_{OCDG} (4xz i^+ + y^2 j^+ + yzk^+) (-i^+) dy dz = \int_0^1 \int_0^1 4(1) z dy dz = 0 \quad (\text{as } x =$$

on putting these values in (1), we get

$$\oint_s F^- \cdot n^{\hat{}} ds = 0 + 1/2 + 6 - 1 + 2 + 0 \Rightarrow 3/2$$

Q 6 (a) Evaluate  $\frac{\partial}{\partial \theta} \{ A^* \{ B * C \} \}$

$$i^{\hat{}} \quad j^{\hat{}} \quad k^{\hat{}}$$

$$(B * C) = \begin{matrix} \cos q & -\sin q & -3 \\ 2 & 3 & -1 \end{matrix}$$

$$= i^{\hat{}} (\sin q + 9) - j^{\hat{}} (-\cos q + 6) + k^{\hat{}} (3\cos q + 2\sin q)$$

$$A^*( B * C ) = \begin{matrix} i^{\hat{}} & j^{\hat{}} & k^{\hat{}} \\ \sin q & \cos q & Q \\ (\sin q + 9) & (\cos q - 6) & (3\cos q + 2\sin q) \end{matrix}$$

$$= i^{\hat{}} [\cos q(3\cos q + 2\sin q) - q(\cos q - 6)]$$

$$- j^{\hat{}} [\sin q(3\cos q + 2\sin q) - q(\sin q + 9)]$$

$$+ k^{\hat{}} [\sin q(\cos q - 6) - \cos q (\sin q + 9)]$$

at  $q = 0$

$$\Rightarrow i^{\hat{}} [\cos 0(3\cos 0 + 2\sin 0) - 0(\cos 0 - 6)]$$

$$- j^{\hat{}} [\sin 0(3\cos 0 + 2\sin 0) - 0(\sin 0 + 9)]$$

$$- j^{\hat{}} [\sin 0(3\cos 0 + 2\sin 0) - 0(\sin 0 + 9)]$$

$$\Rightarrow 3\mathbf{i} - 9\mathbf{k}$$

ans

**Q 6 A particle moves along the curve  $x=t^3 + 1$ ,  $y=t^2$ ,  $z=2t+5$  where  $t$  is the time. Find the components of the velocity and acceleration at  $t=1$  in the direction  $\mathbf{i}+\mathbf{j}+3\mathbf{k}$**

$$x = t^3 + 1, y = t^2, z = 2t + 5$$

$$\mathbf{r} = xi + yj + zk$$

$$\mathbf{r} = (t^3 + 1)\mathbf{i} + (t^2)\mathbf{j} + (2t + 5)\mathbf{k}$$

$$\text{velocity} = \frac{dr}{dt} = 3t^2 \mathbf{i} + 2t \mathbf{j} + 2 \mathbf{k}$$

$$\text{when } t = 1, \text{ we have, } \frac{dr}{dt} = 3 \mathbf{i} + 2 \mathbf{j} + 2 \mathbf{k}$$

$$\text{unit velocity along } (\mathbf{i} + \mathbf{j} + 3\mathbf{k}) = (\mathbf{i} + \mathbf{j} + 3\mathbf{k}) / \sqrt{1+1+9}$$

$$= \frac{1}{\sqrt{11}}(\mathbf{i} + \mathbf{j} + 3\mathbf{k})$$

$$\text{component of velocity } (3 \mathbf{i} + 2 \mathbf{j} + 2 \mathbf{k}) \text{ along } (\mathbf{i} + \mathbf{j} + 3\mathbf{k})$$

$$= (3 \mathbf{i} + 2 \mathbf{j} + 2 \mathbf{k}) \cdot \frac{1}{\sqrt{11}}(\mathbf{i} + \mathbf{j} + 3\mathbf{k})$$

$$= \frac{1}{\sqrt{11}}(3+2+6) = \frac{11}{\sqrt{11}} = \mathbf{11} \text{ ans}$$

## **Q7 (a) Solve by the method of variation of parameter**

$$\frac{d^2y}{dx^2} + n^2y = \sec nx$$

solution: -  $(D^2 + n^2)y = \sec nx$

$$m^2 + n^2 = 0 \Rightarrow m = \pm ni$$

$$cf = c_1 \cos nx + c_2 \sin nx$$

$$y = A' \cos nx + B' \sin nx \quad \text{-----(I)}$$

by diff. Equation (1)

$$(-nA' \sin nx + B' n \cos nx = \sec nx) \quad \text{-----(II)}$$

Now multiplying by  $n \sin nx$  in equation (I) & multiplying by equation (II)

$$n \sin nx (A' \cos nx + B' \sin nx = 0)$$

$$\cos nx (-nA' \sin nx + B' n \cos nx = \sec nx)$$

$$A' n \sin nx \cos nx + B' n \sin^2 nx = 0$$

$$- A' n \sin nx \cos nx + B' n \cos^2 nx = 1$$

---

$$nB' (\sin^2 nx + \cos^2 nx) = 1$$

$$B' = \frac{1}{n}$$

$$\frac{dB}{dx} = \frac{1}{n}$$

$$dB = \frac{dx}{n}$$

$$B = \frac{x}{n} + C$$

$$A'\cos nx + B'\sin nx = 0$$

$$A'\cos nx + \frac{1}{n}\sin nx = 0$$

$$A' = -\frac{1}{n} \frac{\sin nx}{\cos nx} = -\frac{1}{n} \tan nx$$

$$\int dA = -\frac{1}{n} \int \tan nx dx + C_2$$

$$A = -\frac{1}{n^2} \log \sec nx + C_2$$

$$y = \left[ -\frac{1}{n^2} \log \sec nx + C_2 \right] \cos nx + \left( \frac{x}{n} + C_1 \right) \sin nx$$

$$Y = C_1 \sin nx + C_2 \cos nx + \frac{x}{n} \sin nx - \frac{1}{n^2} \cos nx \cdot \log \sec nx$$

**Q 7 (b) solve  $\frac{d^2y}{dx^2} - 2 \tan x \frac{dy}{dx} + 5y = \sec x \cdot e^x$**

$$\frac{d^2y}{dx^2} - 2 \tan x \frac{dy}{dx} + 5y = \sin x \cdot e^x$$

$$y'' - 2 \tan x y' + 5y = \sin x \cdot e^x$$

$$\text{compare with } y'' + 8y' + qy = R$$

$$p = -2 \tan x, q = 6, R = \sin x \cdot e^x$$

$$\text{for C.F} \quad \mu = e^{-\int \frac{1}{2} p dx}$$

$$q_1 = q - \frac{1}{2} \frac{dp}{dx} - \frac{p^2}{4}$$

$$R_1 = \frac{R}{4}$$

$$\mu = e^{-\int \frac{1}{2} -2 \tan x dx} = e^{\log \sec x} = \sec x$$

$$q_1 = 5 \frac{d}{dx}(-2 \tan x) - \frac{4 \tan^2 x}{4}$$

$$= 5 + \sec^2 x - \tan^2 x = 6$$

$$R_1 = \frac{\sin x e^x}{4}$$

$$\frac{d^2v}{dx^2} + q_1 v = R_1$$

$$(D^2 + 6)v = \frac{\sin x e^x}{4}$$

$$A.E = m^2 + 6 = 0 \Rightarrow m^2 = -6 \Rightarrow m = \pm 6i$$

$$CF = (C_1 \cos 6x + C_2 \sin 6x)$$

$$P.I = \frac{1}{(D^2 + 6)} \frac{\sin x e^x}{4} \Rightarrow \frac{1}{4(D^2 + 6)} (\sin x \cdot e^x)$$

$$= \frac{e^x}{4[(D+1)^2 + 6]} \sin x$$

$$= \frac{e^x}{4(D^2 + 1 + 2D + 6)} \sin x \Rightarrow \frac{e^x}{4(-1^2 + 1 + 2D + 6)} \sin x$$

$$= \frac{e^x}{4(-1+1+2D+6)} \sin x \Rightarrow \frac{e^x}{4(2D+6)} \sin x$$

$$= \frac{e^x}{8(D^2-9)} (D-3) \sin x \Rightarrow \frac{e^x}{8(-1-9)} (D-3) \sin x$$

$$= \frac{e^x}{-80} (D-3) \sin x \Rightarrow \frac{e^x}{-80} (D \sin x - 3 \sin x)$$

$$= \frac{e^x}{-80} (\cos x - 3 \sin x)$$

$$\text{C.S} = (C_1 \cos 6x + C_2 \sin 6x) - \frac{e^x}{80} (\cos x - 3 \sin x)$$



Visit [www.goseeko.com](http://www.goseeko.com) to access free study material as per your university syllabus